

DYE DEGRADATION EFFECT OF AN INTERGENERIC HYBRID BETWEEN *GANODERMA LUCIDUM* AND *PLEUROTUS FLORIDA*



Usha Raja Nanthini, A., Mari Selvam, V. and Ranjit Singh, A.J.A*.

¹Department of Zoology, Sri Paramakalyani College, Alwarkurichi, Tamilnadu, India.

*Email: ranjithspkc@gmail.com

Received on: 10 October 2013, accepted on: 12 December 2013

Abstract: Synthetic dyestuffs present a challenge to conventional physico-chemical and biological treatment methods. The study focus on the decolorization of three different textile dyes (Reactive Red 2, Reactive Black 5 and Reactive Yellow 44) by a hybrid mushroom attained by somatic hybridization between an edible mushroom *Pleurotus florida* and a medicinal mushroom *Ganoderma lucidum* and the effect was compared with that of the parents. Maximum amount of protoplasts were released when 5 days old mycelial culture was agitated with 15 mg of lytic enzyme in an osmotic stabilizer (0.6 M $MgSO_4 \cdot 7H_2O$ in 0.05 M Sodium Maleate buffer pH 5) for 18 hrs. After the NTG treatment and mycelial regeneration, arg⁻ and phe⁻ auxotrophic mutants of *Ganoderma lucidum* and *Pleurotus florida* respectively were selected as fusants. A hybrids was obtained when the fusants were incubated in 40% PEG succeeded by screening. The hybrid showed higher rate of decolorization for Reactive Black 5 and Reactive Yellow 44 and rapid decolorization in Reactive Red 2, than the parent strains.

Key words: *Ganoderma lucidum*, *Pleurotus florida*, Protoplast fusion, Reactive dyes degradation.

INTRODUCTION

Synthetic dyes are extensively used in several industries including textile, paper, printing, cosmetics and pharmaceuticals (Marmion, 1991). The total world colorant production is estimated to be 8, 00,000 tons per year and atleast 10% of the used dye stuff enter the environment through wastes (Palmieri *et al.*, 2005; Levin *et al.*, 2004). Decolourisation of textile dye effluent does not occur when treated aerobically by municipal sewage systems (Willmott *et al.*, 1998).

Colour can be removed from effluent by chemical and physical methods including absorption, coagulation- flocculation, ion exchange oxidation and electrochemical methods (Lin *et al.*, 1994). The above ways for clean up are expensive, which limit their application (Moreira *et al.*, 2000). Alternatively, dye decolourisation using microbial enzymes has received great attention in recent years due to its efficient application (Abadulla *et al.*, 2000). A number of biotechnological approaches have been suggested by recent research as of potential interest towards combating pollution by dyes in an ecofriendly manner. They include the use

of bacteria and fungi or often in combination with physico- chemical processes (Willmott *et al.*, 1998). Though bacteria could utilize dyestuff under anoxic conditions, the disadvantage of this method is the production of aromatic amines by these organisms. These amines may be toxic and carcinogenic (Meyer, 1981). Aerobic bacteria usually tend to be specific towards a particular dye (Reddy, 1995).

White rot fungi are unique among eukaryotic and prokaryotic microorganisms, because they possess a very powerful extra cellular oxidative enzymatic system (LDS) (Rigas and Drista, 2006). Their extracellular ligninolytic system involve enzymes such as Manganese peroxidases (MnP), Lignin peroxidases (LiP) and Laccases (Lac), which are involved not only in the degradation of lignin, but also in the degradation of various xenobiotic pesticides, PCBs and PAHs (Pointing, 2001; Barr and Aust, 1994; Kubatova *et al.*, 2001; Wesenburg *et al.*, 2003; Jarosz-Wilkolazka *et al.*, 2002; Kirk and Farrell, 1987). They being non-specific can attack a wide variety of complex aromatic dyestuffs (Barclay *et al.*, 1990; Nagai *et al.*, 2002; Boer *et al.*, 2004; Kamitsuji *et al.*, 2005)

Since the enzymes are extracellular, the substrate distribution limitation into the cell generally encountered in bacteria, is not observed. Further, the extracellular enzyme system enables white rot fungi to tolerate high pollutant concentration (Kapdan *et al.*, 2000; Rodriguez *et al.*, 2004).

Revankar and Lele (2006) have reported that a new isolate of white rot fungus, *Ganoderma* sp. WR-1 showing very high laccase producing ability. Asgher *et al.* (2006) has reported decolorization of some reactive textile dyes by *Ganodemalucidum* and *Pleurotosteatatus*, Yet there is no reports on decolorization using fungal hybrids.

Mushrooms are an abundant source of a wide range of useful natural products with biological activities. Protoplasts are now regarded as a useful tool for genetic manipulations (Peberdy, 1980; Ferenczy, 1981), which involve fusion leading to somatic hybridization and gene transformation. As conventional hybridization, protoplast fusion can be performed intraspecifically (Kiguchi and Yanagi, 1985; Toyomatsu and Mori, 1987), interspecifically (Takehara *et al.*, 1993; Matsumoto *et al.*, 1997), inter-generically (Eguchi *et al.*, 1993; Zhao and Chang, 1996) and even inter-heterogenerically (Eguchi and Higaki, 1995; Toyomatsu and Mori, 1987). *Ganoderma* is a polypore mushroom that is soft, corky and flat. The mushroom is too tough to be edible (Wasser, 2005). It has been an economically imported species, particularly in the Far East countries (china, Japan, Korea, etc.,) for over 4000 years. It is widely grown on a commercial scale and is commonly purchased for its medicinal spiritual properties. *Pleurotus florida*, commonly known as oyster fungus, grows wildy in tropical and subtropical rainforests, and can be artificially cultivated. This mushroom is considered as a delicacy and treasured for its flavor and taste. They are rich in nutrition and are claimed to have medicinal value (Zadrazil, 1980). A step has been taken to construct a somatic hybrid which may show recombined characteristics of both *Ganoderma* and *Pleurotus*. A few reports are available on the development of regenerating mushroom protoplasts, but usage of macerozyme, and selection of auxotrophic mutants Arg and Phe

has not been reported. The present study focuses on producing a somatic hybrid between *Ganodemalucidum* and *Pleurotus florida* through protoplast fusion technology and finding its effect on decolorizing three reactive dyes and comparing it with the effect of parents.

MATERIALS AND METHODS

Reactive dyestuffs

The three reactive textile dyestuffs were Reactive Red 2 (Red M₂ B), Reactive, Reactive Black 5 (Black B) and Reactive Yellow 44 (Yellow MR), provided by the distribution office at Pallipalayam, Erode, India.

Strain and culture

A mycelial culture of *Pleurotus florida* was developed from a local commercial strain and pure culture of *Ganoderama lucidum* (MTCC) were used. Both the strains were cultured in Potato Dextrose broth and Malt Dextrose broth. Mycelia that had grown for 5 days were used for protoplast formation. The cultures were subcultured periodically.

Culture media

The basal media contained (g/l of distilled water), Starch 20, Yeast extract 2.5; KH₂PO₄ 1.0; NaH₂PO₄ 0.05; MgSO₄·7H₂O 0.5; CaCl₂ 0.01; FeSO₄·7H₂O 0.01; MnSO₄·4H₂O 0.001; CuSO₄·5H₂O 0.002. The pH of the medium was adjusted to 5.5 (Revanker and Lele, 2006).

Isolation of Protoplast

Four day-old monokaryotic mycelia of *Ganodermalucidum* and *Pleurotus florida* was transferred onto 100ml of Malt extract broth plus 20 glass beads (0.50 cm in diameter) with the agitation speed of 200 rpm, at 25°C for 4 days. Protoplasts were separated from the mycelia using a method modified from Hashiba (1992) by transferring 0.3 g of the mycelia onto 3 ml of the sterilized lytic enzyme solution having macerozyme in a test tube and shaking it at 100 rpm at room temperature for 16 hrs.

Effect of lytic enzyme

Macerozyme was tested for their effect on protoplast release in *Ganoderma lucidum* and *Pleurotus florida*. Macerozyme was used at

various concentrations of 8mg, 10mg, and 15mg to digest mycelium. It was prepared in 0.6mol/l sodium Maleate buffer(pH5.8) along with MgSo₄.

Effect of incubation time:

Effect of incubation time in release of protoplast in *Ganoderma lucidum* and *Pleurotus florida* when incubated with macerozyme was analyzed for various periods of 2hr, 10hr, 16hr, and 18hr.

Effect of Inoculum size

Freeze dried mycelia with different weight was tested for their effect in release of protoplast. Mycelia were taken in different dry weight (0.25g,0.5g,1g and 1.5g) in addition to osmotic stabilizer and lytic enzyme.

Purification of Protoplast

The mycelial remnants were removed by filtration through a sintered glass filter and the suspended protoplasts were precipitated at 1000xg for 10 min. The protoplasts were then washed twice with the osmotic stabilizer in 1 ml of the osmotic stabilizer which contained 0.6 M MgSO₄·7H₂O in 0.05 M Sodium Maleate buffer, pH 5. And they were finally suspended in 5 ml of the osmotic stabilizer. Protoplasts obtained from *Ganoderma lucidum* and *Pleurotus florida* were counted using a haemocytometer.

Mutagenic procedure

Protoplast suspension (ca. 1.2 x 10⁶/ml) was treated with N-methyl-N'-nitro-N-nitrosoguanidine (NTG) for mutagenesis. The protoplasts were exposed to the action of NTG for varied period of time from 15 to 120 seconds. NTG was used at two concentrations, 15 and 30µg/ml in osmotic stabilizer solution.

Regeneration

The protoplasts were immediately washed thrice with osmotic stabilizer to remove NTG and diluted to 1x10⁴ protoplasts/ml. 0.1 ml from the dilution was used for protoplast regeneration on a plate with regeneration medium. The plate was incubated at 25°C until colonies occurred. Each colony was isolated and inoculated in Malt extract agar slants.

Selection of mutants

To select the mutants, all the colonies from regenerating medium were transferred to minimal medium (MM) without amino acids. Regenerated colonies those did not grow on MM were tested for their requirements by further transfer onto MM containing different amino acids (20 µg/ml). Genotypes were considered to be stable after five successive subcultures on appropriately supplemented MM.

Protoplast Fusion

Freshly prepared protoplasts (diluted to be 1x10⁶ protoplasts/ml) of *Ganoderma lucidum* and *Pleurotus fusion* were mixed and centrifuged at 1000x g for 10 min. The supernatant was rinsed off and 1 ml of sterilized PEG (6000) (40 g PEG in 100 ml 0.05 M CaCl₂·2H₂O) was added to the mixture and incubated at room temperature for 20 minutes. The tubes were intermittently shaken for every 5 min and then centrifuged at 1000x g for 10 minutes. The supernatant was discarded and the protoplast mixture was taken for hybrid screening.

Decolourisation procedure

Decolorization flasks were prepared in triplicates each containing 50 ml of nutrient media for three days observation. The medium was sterilized (121°C) in an autoclave for 15 min. Each flask was inoculated with 5 numbers of agar plug cut from PDA plates, and allowed to grow for four days in 30°C. Fourth day three dye stuffs were added in a concentration of 100 mg/l separately in 3 sets. Biotic control (with inoculum and without dye) and abiotic control (with dyestuff and nutrient, but received no inoculum) were maintained. For every 24 hours samples were removed, centrifuged at 8,000x g for 20 minutes and the supernatant obtained was used for the determination of residual dyestuff concentration.

Dyestuff analysis

Absorbance measurements were done with UV/Visible spectrometer. When the wavelength with maximum absorbance (λ_{max}) was measured, it was 440, 590 and 420 nm for the dyes Reactive Red 2, Reactive Black 5 and Reactive Yellow 44 respectively. Respective blanks (containing only medium, omitting the

respective dye) were used to calculate the absorbance values for supernatants collected at time intervals. The absorbance values were used to know the percentage of decolorization.

Decolorisation% =

$$\frac{\text{Initial absorbance} - \text{observed absorbance}}{\text{Initial absorbance}} \times 100$$

RESULTS AND DISCUSSION

The study is based on the formation, fusion, regeneration of *Ganoderma lucidum* and *Pleurotus florida* protoplasts and application of the hybrid in textile dye decolorization.

Isolation of Protoplast

The commercially prepared enzymes Novozyme 234 and Cellulase CP have been used previously to effect protoplast release from a variety of fungal species (Hamlyn *et al.*, 1981). Our results extend the use of macerozyme enzyme to release protoplasts from *Ganoderma lucidum* and *Pleurotus florida*. Although pretreatment with thiols was required for efficient protoplast release in some fungal agents (Brown, 1971; Fincham and Day, 1971.), no such pretreatment was necessary for *Ganoderma lucidum* and *Pleurotus florida*. The amount of protoplasts obtained in *Ganoderma lucidum* and *Pleurotus florida* were 3×10^5 and 1.8×10^5 protoplasts/ml, respectively. The results achieved correlate to some extent with those published by Dhitaphichit and Pornsuriya (2005).

Production of protoplast, however, varies with the factors used in the isolation process e.g. species and age of fungal mycelia, type and condition of the lytic enzyme (Peberdy, 1989) and of the osmotic stabilizer (Peberdy and Fox, 1993) and complex interactions exist between them.

Effect of lytic enzyme

Among the various concentrations (8mg, 10mg, and 15mg) of lytic enzyme Macerozyme used, both the strains showed more yield in 15 mg/ml concentration (Fig. 1).

Effect of incubation time

Release of protoplasts was found to begin after 10h of incubation and attained maximum at 18h (Fig. 2).

Effect of Inoculum size

When freeze dried mycelia were taken in different weight (0.25g, 0.5g, 1g and 1.5g), the highest percentage of protoplast release was observed in 1.5g of inoculum (Fig. 3).

Effect of osmotic stabilizer

Protoplasts released from *Ganoderma lucidum* and *Pleurotus florida* were found to be osmotically fragile spherical bodies, mostly of uniform size, accompanied by a few relatively bigger ones. Among different concentrations of lysing enzymes tried, we optimized that 15mg/ml with 0.6M MgSo₄ and Sodium Maleate buffer as osmotic stabilizer to release higher number of protoplast from *Ganoderma* and




Fig. 1. Effect of enzyme concentration on protoplast yield




Fig. 2. Effect of incubation time on protoplast yield




Fig. 3. Effect of inoculum size in the release of protoplast

Pleurotus sp. However, Pe'er and Chet (1990) obtained highest protoplasts from *T. harzianum* using Novozym 234 at 10 mg/ ml with 0.6M KCl and Tschien and Li used 15 mg/ ml of Novozym with 0.6M sucrose to isolate maximum protoplasts from *T. harzianum* and *T. Koningii*. Further, Balasubramanian *et al.* (2003) reported maximum number of protoplasts from *Trichothecium roseum* using Novozym 234 in combination with chitinase and cellulase each at 5mg/ml.

Regeneration

In the process of protoplast regeneration, the critical element required for stabilizing is the hypertonic solution. It was observed that the choice of a suitable osmotic stabilizer solution has a pronounced effect on regeneration. Maleic acid - NaOH buffer (pH- 5.0) with 0.6M $MgSO_4$ was the osmotic stabilizer reported for *Schizophyllum commune*, *Coprinus pel lucidus*

and *Coprinus cinereus* (De Varies *et al.*, 1972) and the same was found to be most suitable in the present study too.

Protoplasts that survived and regenerated after various exposures were screened for any auxotrophy in minimal medium with aminoacids. It should be mentioned that although a number of colonies were identified as auxotroph on initial screening, most of them quickly reverted back to the wild type on subsequent subculturing. Among the mutants, those did not revert back for successive five generations were selected for the further study are auxotrophic mutant was identified in *Ganoderma lucidum* and phe⁻ auxotrophic mutant was identified in *Pleurotus florida* (Fig. 7).

Protoplast fusion

The molecular weight of the PEG and concentration of the fusogen employed seem to be critical to obtain optimal fusion. Inter-




Fig. 7 a. arg⁻ Auxotroph- *Ganoderma lucidum*; b. phe⁻ Auxotroph- *Pleurotus florida*




Fig. 8. Regeneration of fused protoplast

generic protoplasts fusion between *Ganoderma lucidum* and *Pleurotus florida* was achieved by employing 40% PEG (600) similar to that reported by Mrinalini *et al.*, (1996). Regeneration of mycelia happened when inoculated on regeneration media (Fig 8).

Hybrid selection:

After regeneration, selection of hybrid was carried out by inoculation on different synthetic media. The hybrids that need either arginine or




Fig. 9. Confirmation of hybrid grown on minimal medium after protoplast fusion (a. minimal medium with Arginine, b. minimal medium with Phenyl alanine, c. minimal medium with arg+ phe, d. growth on minimal medium.

Table 1. Decolorization of Reactive Black 5 (Black B)

Day	Decolorization(%)		
	<i>Pleurotus florida</i>	<i>Ganoderma lucidum</i>	Hybrid
1	0	0	0
3	0	4.4	13.13
4	4.04	5.05	20.2
5	14	8.58	45.5
6	25.75	28.3	44.14
7	46	29.3	59.75
8	51.01	61.25	76.76
9	67.17	74.75	77.78
10	87.37	87.37	94.94




Fig. 4. Decolorization of Reactive Black 5 (Black B)




Fig. 5. Decolorization of Reactive Yellow 44 (Yellow MR)




Fig. 6. Decolorization of reactive red 2 (Red M₅B)

Table 2. Decolorization of Reactive Yellow 44 (Yellow MR)

Day	Decolorization(%)		
	<i>Pleurotusflorida</i>	<i>Ganodermalucidum</i>	Hybrid
1	44.4	55.45	43.3
2	57.8	66.4	57.8
3	66.7	77.8	59.3
4	74.3	78	75
5	74	79	75.5
6	80	79.3	82.22
7	82.2	80	82.22
8	84	80	84.4
9	84	81.2	85.3
10	84.1	82.22	86.66

Table 3. Decolorization of reactive red 2 (Red M₅B)

Day	Decolorization(%)		
	<i>Pleurotusflorida</i>	<i>Ganodermalucidum</i>	Hybrid
1	6.5	2.1	6.5
2	10.8	13	15
3	13	23.9	23.2
4	15.5	34.7	36.5
5	21.7	45.6	54.3
6	32.6	52.3	65
7	49	69.5	73.9
8	65.2	71.7	74.78
9	72	73.91	84.78
10	89.13	74.2	86.9

phenylalanine are intraspecific fusion products and they were eliminated. The mutants that were stable, retaining the auxotrophy after several subcultures, did not require arginine and phenylalanine supplementation in the synthetic medium after fusion were considered as the hybrids (Fig. 9) and used further.

Dye decolorization and Dyestuff analysis:

Results of time course studies on decolorization of Reactive Black 5 (Black B) are displayed in Table 1, Fig. 4. Although the rate of colour removal was nil within the first 3 days for *Pleurotus florida* and 2 days in the case of both the hybrid and *Ganoderma lucidum*, hybrid has brought the highest rate of decolorization (94.94%) than the parent strains. Both *Pleurotus florida* and *Ganoderma lucidum* could bring about 87.37% decolorization after 10 days of incubation.

A time course study on decolorization of Reactive Yellow 44 (Yellow MR) is shown in the Table 2, Fig. 5. Even though Reactive Yellow 44 (Yellow MR) was decolorized effectively upto 84% in 8 days by *Pleurotus florida* and there was no further change till 10 days of incubation. 82.2% of decolorization was observed by *Ganoderma lucidum*. The hybrid showed the highest rate of decolourisation (86.7%) when compared to the parents.

Time course studies on decolorization of reactive red 2 (Red M₅B) is exhibited in the Table 3, Fig. 6. *Ganoderma lucidum* showed the least rate of decolorization (74.2%) and *Pleurotus florida* exhibited the highest rate of decolorization (89.13%) in 10 days. But more than 50% decolorization was obtained only after 7 days of incubation. Eventhough the decolorization rate showed by the hybrid was comparatively lesser than that of *Pleurotus florida*, it pronounced more than 50% decolourisation within 5 days of incubation.

It has been reported that azo dyes are recalcitrant to decolourisation and could only be decolourised to a limited extent (Nyanhongo *et al.*, 2002). But *Ganoderma lucidum* and *Pleurotus florida* had efficiently degrade Reactive red 2 (red M₅ B), Reactive Black 5

(Black B) and Reactive Yellow 44 (Yellow MR). As a highlight the hybrid has shown more efficiency in decolorizing all the dyestuffs than the parent strains.

Physiological differences among *Ganoderma lucidum*, *Pleurotus florida* and the hybrid may account for difference in their decolorization abilities. The complex enzymatic system responsible for dye degradation and pattern of its expression may also vary. Intergeneric hybrid of *Ganoderma lucidum* and *Pleurotus florida* had exhibited good potential for decolorization of relatively less studied reactive textile dyestuffs. It may be due to enhancement in the enzymatic system occurred by fusion.

CONCLUSIONS

The results of the study will form the basis for development of cost effective and robust indigenous technology for bioremediation of these reactive dyes and dye based effluents.

ACKNOWLEDGEMENTS

The authors thank the University Grants Commission (UGC), India for their financial support to carry out this study.

REFERENCES

- Abadulla, E., Tzanov, T., Costa, S., Robra, K.H., Cavaco-Paulo, A., Gubitz, G.M. 2000. Decolorization and detoxification of textile dyes with a laccase from *Trametes hirsuta*. *Applied environmental microbiology*, 66: 3357-62.
- Asgher, M., Shah, S.A.H., Ali, M., Legge, R.L. 2006. Decolorization of some reactive textile dyes by white rot fungi isolated in Pakistan. *World Journal of Microbiology and Biotechnology*, 22: 89-93.
- Balasubramanian, N., Annie Juliet, A., Srikalaivani, P., Lalithakumari, D. 2003. Release and regeneration of protoplasts from *Trichothecium roseum*. *Can. J. Microbiol.*, 49:263-8.
- Barclay, C.D., Moore, D.M., Lander, S.R., Legge, R.L. 1990. Heat denaturation kinetics of lignin peroxidases from *Phanerochaete chrysosporium*. *Enzyme and Microbial Technology*, 12:778-782.

- Barr, D.P., Aust, S.D. 1994. Mechanism white-rot fungi use to degrade pollutants. *Environmental science and technology*, 28: 78-87.
- Boer, C.G., Obici, L., Souza, C.G.M. de., Piralta, R.M. 2004. Decolorization of synthetic dyes by solid state cultures of *Lentinula (Lentinus) edodes* producing manganese peroxidase as the main lignolytic enzyme. *Bioresource Technology*, 94: 107-112.
- Brown J.P. 1971. Susceptibility of the cell walls of some yeast to lysis by enzymes of Helvonomati. *Can. J. Microbiol.*, 17: 205-208.
- De Vries O.M.H., Wessels. J.G.H. 1972. Release of protoplasts from *Schizophyllum commune* by a lytic enzyme preparation from *Trichoderma viride*. *J. Gen. Microbiol.*, 73: 13-22.
- Dhitaphichit, P. and Pornsuriya. C. 2005. Protoplast fusion between *Pleurotostreatus* and *P. djamor Songklanakarini*. *J. Sci. Technol.*, 27(5): 975-982.
- Eguchi, F. and Higaki, M. 1995. Production of new species of edible mushrooms by protoplast fusion and analysis of the fusant between *Pleurotussajorcaju* and *Mycocleptodonoi-desaitchisonii*. *Mokuzai gakkaiishi*, 41(3): 342-348.
- Eguchi, F., Fukuzumi, T. and Higaki, M. 1993. Production of new species of edible mushrooms by protoplast fusion Method II. Analysis of the Mycelial and Basidiospores of a Fusant between *Pleurotostreatus* and *Agrocybecy lindracea*. *Mokuzai Gakkaiishi*, 39(7): 825-830.
- Ferenczy, L. 1981. Microbial protoplast fusion. In *Genetics as a Tool in Microbiology*, ed. Glover, S.W. and Hopwood, D.A. 1-34. Cambridge University Press.
- Fincham, F.R.S. and Day, P.R. 1971. Fungal genetics. Blackwell Scientific Publications. Ltd.. Oxford. England.
- Hamlyn, P.F., Bradshaw, R.E., Mellon, F.M., Santiago, C.M., Wilson, J.M., Peberdy, J.F. 1981. Efficient protoplast isolation from fungi using commercial enzymes. *Enzyme Microb. Technol.*, 3: 321-325.
- Hashiba, T. 1992. Isolation of fungal protoplast. In *Hand book of fungal Mycology*. (Eds: Arora, D.K., Mukerji, K.G. and Marth, E.H) Mercel dekker, Inc., New York. 4: 129-149.
- Jarosz-Wilkolazka, A., Kochmanska-Rdest, J., Malarczyk, E., Wards, W., Leonowicz, A. 2002. Fungi and their ability to decolorize azo and anthraquinonic dyes. *Enzyme microbial technology*, 30: 566-72.
- Kamisuji, H.Y., Watanabe, T., and Kuwahara, M, 2005. Mn²⁺ is dispensable for the production of active MnP₂ by *Pleurotostreatus*. *Biochemical and Biophysical Research Communications*, 327: 871-876.
- Kapdan, K.I., Karki, F.G. M. cullan, G. and Merchant, R. 2000. Biological decolorization of textile dye stuff by *coriolus versicolor* in a packed column reactor. *Environment Technology*, 21: 231-236.
- Kiguchi, T. and Yanagi, S.O. 1985. Intraspecific Heterokaryon and fruit body formation in *Coprinus macrorhizus* by protoplast fusion of Auxotrophic mutants. *Appl. Microbiol. And Biotechnol.*, 22: 121-127.
- Kirk, T.K. and Farrell, R.L. 1987. Enzymatic "Combustion". The microbial degradation of lignin. *Annual Review on Microbiology*, 41: 465-505.
- Kubatova, A, Erbanova L., Eichlerova, L. Homolka, L., Nerud, F., Sasek, V. 2001. PCB congener selective biodegradation by the white-rot fungus *Pleurotostreatus* in contaminated soil. *Chemosphere*, 43: 207-15.
- Levin, L, Papinutti, L. and Forchiassin, F. 2004. Evaluation of argentine white rot fungi for their ability to produce lignin modifying enzymes and decolorize industrial dyes. *Bioresource Technology*, 94 : 169-176.
- Lin, S.H. and Peng, F.C. 1994. Treatment of textile waste water by electrochemical methods. *Water research*, 28: 277-82.
- Marmion, D.M. 1991. *Hand book of US colorants foods, drugs, cosmetics and medical devices*. 3rded, New York: Wiley.
- Matsumoto, M., Eguchi, F. and Higaki, M. 1997. Polyethylene Glycol Centrifugal Fusion between *Pleurotostreatus* and *Pleurotussajorcaju*. *Mokuzai Gakkaiishi*, 43(2):215-219.

- Meyer, U. 1981. Biodegradation of synthetic organic colorant. *FEMS Symposium*, 12: 371-385.
- Moreira, M.T., Mielgo, I., Feijoo, G. and Lema, J.M. 2000. Evaluation of different fungal strains in the decolorization of synthetic dyes. *Biotechnology letters*, 22: 1499-503.
- Mrinalini, C. and Lalithakumari, D. 1996. Protoplast fusion: a biotechnological tool for strain improvement of *Trichoderma* spp. In: Recent development in biocontrol of plant pathogens. New Delhi, India: Today and Tomorrow Printers and Publishers; 133-46.
- Nagai, M., Sato, T., Watanabe, H., Saito, K., Kawata, M. and Enei, H. 2002. Purification and characterization of an extracellular laccase from the edible mushroom *Lentinula edodes* and decolorization of chemically different dyes. *Applied Microbiology and Biotechnology*, 60: 327-335.
- Nyanhongo, G.S., Gomes, J., Gubitz, G.M., Zuauya, R.M., Read, J.S., Steiner, W. 2002. Decolorization of textile dyes by laccase from a newly isolated strain of *Trametes modesta*. *Water res.*, 36: 1449-1456.
- Palmieri, G., Cennamo, G., Sannia, G. 2005. Remazol Brilliant R decolorization by the fungus *Pleurotus ostreatus* and its oxidative enzymatic system. *Enzyme and Microbial Technology*, 36: 17-27.
- Pe'er, S. and Chet, I. 1990. *Trichoderma* protoplast fusion; a tool for improving biocontrol agents. *Can. J. Microbiol.*, 36: 6-9.
- Peberdy, J.F. and Fox, H.M. 1993. Protoplast technology and edible mushrooms. In: Genetics and breeding of mushroom, 125-155 (eds: Chang, S.T., Buswell, J.A., Miles, P.G.). Gordon and Breach Science Publishers.
- Peberdy, J.F. 1980. Protoplast fusion: a tool for genetic manipulation and breeding in industrial microorganisms. *Enzyme Microbiol Technol.*, 2: 23-9.
- Peberdy, J.F. 1989. Fungi without coats—protoplasts as tools for mycological research. *Mycol Res.*, 93: 1-20.
- Pointing, S.B. 2001. Feasibility of bioremediation by white-rot fungi. *Applied Microbiology and Biotechnology*, 57: 20-33.
- Reddy, C.A. 1995. The potential for white rot fungi in the treatment of pollutants. *Current Opin Biotechnology*, 6: 320-328.
- Revankar, M.S. and Lele, S.S. 2006. Enhanced production of laccase using a new isolate of white rot fungus WR-1. *Proc. Biochem.*, 41(3): 581-588.
- Rigas, F. and Dritsa, V. 2006. Decolorization of a polymeric dye by selected fungal strains in liquid cultures. *Enzyme and Microbial Technology*, 39: 120-124.
- Rodriguez Couto, S., Sanroman, M.A., Hofer, D. and Gubitz, G.M. 2004. Stainless steel sponge: a novel carrier for the white rot fungus *Trametes hirsuta* for decolorization of textile dyes. *Bioresource Technology*, 95: 67-72.
- Takehara, T., Kumata, A. and Aono, S. 1993. Inter-specific protoplast fusion between some *Pleurotus* species using auxotrophic mutants. *Mokuzai Gakkaishi*, 39(7): 855-859.
- Toyamatsu, T. and Mori, K.I. 1987. Intra- and inter-specific protoplast fusion between some *Pleurotus* species. *Agric. Biol. Chem.*, 51 (3): 935-937.
- Wasser, S.P. 2005. Reishi or Ling Zhi (*Ganoderma lucidum*) Encyclopedia of Dietary Supplements. Marcel Dekker, Inc., 603-622.
- Wesenberg, D., Kyriakides, I. and Agathos, A. 2003. White rot fungi and their enzymes for the treatment of industrial dye effluents. *Biotechnology advanced*, 22: 161-187.
- Willmott, N., Guthrie, J. and Nelson, G. 1998. The biotechnology approach to color removal from textile effluent. *JSDC*, 114: 38-41.
- Zadrazil, F. 1980. Influence of ammonium nitrate and organic supplements on the yield of *Pleurotus sajor-caju* (Fr.) Sing. *European J. Appl. Microbiol. Biotechnol.*, 9: 31-35.
- Zhao, J. and Chang, S.T. 1996. Intergeneric Hybridization between *Pleurotus ostreatus* and *Schizophyllum commune* by PEG-induced protoplast fusion and Electrofusion. *World J. Microbiol. and Biotechnol.*, 12: 573-578.